

A WIDE-RANGING AND AUTHORITATIVE GUIDE
CONTAINING OVER 6,250 ENTRIES



Oxford

A horizontal decorative bar consisting of a red square on the left, a dark blue square in the middle, and a red square on the right. In the center of the dark blue square is a black and white photograph of a cracked, dry earth surface.

DICTIONARY OF

Earth Sciences

MICHAEL ALLABY

A Dictionary of

Earth Sciences

THIRD EDITION

Edited by

MICHAEL ALLABY

OXFORD
UNIVERSITY PRESS

OXFORD

UNIVERSITY PRESS

Great Clarendon Street, Oxford OX2 6DP

Oxford University Press is a department of the University of Oxford.
It furthers the University's objective of excellence in research, scholarship,
and education by publishing worldwide in

Oxford New York

Auckland Cape Town Dar es Salaam Hong Kong Karachi

Kuala Lumpur Madrid Melbourne Mexico City Nairobi

New Delhi Shanghai Taipei Toronto

With offices in

Argentina Austria Brazil Chile Czech Republic France Greece

Guatemala Hungary Italy Japan Poland Portugal Singapore

South Korea Switzerland Thailand Turkey Ukraine Vietnam

Oxford is a registered trademark of Oxford University Press
in the UK and in certain other countries

Published in the United States

by Oxford University Press Inc., New York

© Oxford University Press 1990, 1999, 2003, 2008

The moral rights of the authors have been asserted

Database right Oxford University Press (maker)

First published 1990

First issued as an Oxford University Press paperback 1991

Second edition 1999

Reissued with new covers and corrections 2003

Third edition 2008

All rights reserved. No part of this publication may be reproduced,
stored in a retrieval system, or transmitted, in any form or by any means,
without the prior permission in writing of Oxford University Press,
or as expressly permitted by law, or under terms agreed with the appropriate
reprographics rights organization. Enquiries concerning reproduction
outside the scope of the above should be sent to the Rights Department,
Oxford University Press, at the address above

You must not circulate this book in any other binding or cover
and you must impose this same condition on any acquirer

British Library Cataloguing in Publication Data

Data available

Library of Congress Cataloguing in Publication Data

Data available

Typeset by SPI Publisher Services, Pondicherry, India

Printed in Great Britain

on acid-free paper by

Clays Ltd, St Ives plc

ISBN 978-0-19-921194-4

1 3 5 7 9 10 8 6 4 2

From the Preface to the First Edition

Setting intellectual boundaries that would serve to define ‘geology’ has never been simple. As long ago as 1830, in his *Principles of Geology*, Charles Lyell expressed the view that geologists should be well versed in chemistry, natural philosophy, mineralogy, zoology, comparative anatomy, and botany. For at least a century and a half those who would study the structure and composition of the Earth have had to familiarize themselves with a wide range of scientific disciplines.

Strictly speaking, the word ‘geology’ describes all studies of the Earth. Traditionally, however, ‘geology’ has come to mean the study of rocks.

T. C. Chamberlin used the name ‘Earth sciences’ to embrace astronomy, cosmogony, and cosmology as well as the traditional disciplines, and Alfred Wegener (originally a meteorologist) also used it, but it was not until the 1960s that it began to gain a wider currency. Within ten years it was widely accepted, used sometimes in the singular, nowadays commonly in the plural. When, in the late summer of 1985, our friends at the Oxford University Press invited us to compile a dictionary of terms used in the topics directly related to studies of the Earth, it was clear that it should be a dictionary of ‘Earth sciences’.

We had to begin by defining the term for our own purpose. We examined the way it was used by other authors, assembled a kind of consensus, and determined that our dictionary should include terms from climatology, meteorology, economic geology, engineering geology, geochemistry, geochronology, geomorphology, geophysics, hydrology, mineralogy, oceanography, palaeoclimatology, palaeoecology, palaeogeography, palaeontology, pedology, petrology, the philosophy and history of the Earth sciences including brief biographical notes of important figures, planetary geology, sedimentology, stratigraphy, structural geology, tectonics, and volcanology.

The task of a dictionary is descriptive, not prescriptive. It records words and expressions that are in current use and explains the meanings attached to them, but it does not impose those meanings or seek to dictate what a correct usage should be. As recorders, we express no opinions.

We would emphasize that the book is meant to be used as a dictionary. In no sense is it intended to be a textbook in its own right.

Preface to the Second Edition

Work on the first edition of *The Concise Oxford Dictionary of Earth Sciences* was completed in the summer of 1988, and the book was published in 1990. Many changes and advances have occurred during the years that have passed since the initial compilation and we are glad to have been given the opportunity to take account of them in a second edition.

The revision has been extremely thorough. Every entry from the first edition has been scrutinized and many have been amended to bring them up to date.

In preparing the new edition we have been assisted by Dr Robin Allaby. His contribution has greatly strengthened the revision and we are very grateful for his help.

We also would like to thank Professor D. H. Tarling and Dr C. D. Gribble. They each scrutinized a long list of entries, revising them where necessary.

Finally, we thank Nigel May, the science librarian at the library of the University of Plymouth, for allowing us to make use of the library facilities.

AILSA ALLABY
MICHAEL ALLABY

Preface to the Third Edition

The second edition of Oxford's *Dictionary of Earth Sciences* was published eight years ago, in 1999, and although the dictionary was reprinted with corrections in 2003, it was possible then to make only minor changes. This edition is entirely new. Every entry has been scrutinized and brought up to date where necessary. New entries have been added and the number of diagrams has increased.

The new entries added in the second edition related predominantly to the expansion of planetary exploration and satellite technology. Those entries have been updated. Many of the new entries in this edition serve to expand the dictionary's coverage of geomorphological terms.

It is in the nature of dictionaries to grow longer with each revision, and I make no apology for the fact that this one conforms to that rule. Such growth is unavoidable, because the language itself is always growing, but when new terms are introduced it does not mean that older terms are discarded. Even terms that do become obsolete do so gradually, surviving for many years in established and much loved texts.

The first edition was compiled with the help of many contributors and advisers, and a smaller number of contributors assisted in preparing the second edition. The value of their hard work endures, and I gladly and fully acknowledge it, for without it there would have been no dictionary to revise. I have worked alone in preparing this third edition, however, so if any new errors have crept in they are mine alone.

MICHAEL ALLABY
TIGHNABRUAICH, ARGYLL
www.michaelallaby.com

Contents

<i>Contributors and Advisers</i>	xi
A Dictionary of Earth Sciences	1
APPENDIX A: Stratigraphic Units as Defined in the North American Stratigraphic Code, 1983	635
APPENDIX B: Time-Scales	637
APPENDIX C: Wind Strength	642
APPENDIX D: SI Units, Conversions, and Multiples	644
APPENDIX E: List of useful websites	646
<i>Bibliography</i>	647

Contributors and Advisers

Ailsa Allaby

Michael Allaby

Robin Allaby, University of Manchester Institute of Science and Technology

Dr Keith Atkinson, Camborne School of Mines

Dr R. L. Atkinson, Camborne School of Mines

Dr T. C. Atkinson, University of East Anglia

Dr A. V. Bromley, Camborne School of Mines

Denise Crook

J. G. Cruickshank, Department of Agriculture for Northern Ireland, Belfast

Dr P. Francis, Open University; Lunar and Planetary Institute, Houston

Professor K. J. Gregory, Goldsmith's College, University of London

Dr C. D. Gribble, University of Glasgow

Dr Colin Groves,* Australian National University

Dr W. J. R. Harries, University of Plymouth

Professor M. Hart, University of Plymouth

Professor Emeritus H. H. Lamb,* University of East Anglia

John Macadam

Dr R. J. T. Moody, Kingston University

Dr J. Penn, Kingston University

Dr John M. Reynolds, Reynolds Geo-Science Ltd.

Dr D. Rolls, Kingston University

Dr I. Roxburgh

Dr N. A. Rupke, Wolfson College, Oxford

Dr Stuart Scott, University of Plymouth

Dr B. W. Sellwood, University of Reading

Dr P. J. C. Sutcliffe, Kingston University

Professor D. H. Tarling, University of Plymouth

Joan Taylor

Professor S. R. Taylor, Australian National University

Dr R. J. Towse,* Kingston University

Dr I. Tunbridge, University of Plymouth

Dr C. E. Vincent, University of East Anglia

Professor Brian F. Windley, University of Leicester

Andrew Yelland, Birkbeck College, London

* Contributor to *The Oxford Dictionary of Natural History* whose earlier entries have been transferred to this book.



aa See LAVA.

AABW See ANTARCTIC BOTTOM WATER.

AAC See ANTARCTIC CONVERGENCE.

Aalenian A *stage in the European Middle *Jurassic (175.6–171.6 Ma, Int. Commission on Stratigraphy, 2004). See also DOGGER.

AAV See AGGREGATE TESTS.

Ab See ALKALI FELDSPAR.

abandoned channel A former stream channel through which water no longer flows (e.g. a *cut-off).

abandonment facies association A *facies association formed under conditions of rising sea level, when *clastic deposition has ceased and sediment is deposited very slowly.

abapical A directional term meaning away from the shell *apex.

abaptation The process by which an organism is fitted to its environment as a consequence of the characters it inherits, which have been filtered by *natural selection in previous environments. Because present environments seldom differ greatly from recent past environments, adaptive fitness can resemble *adaptation. In this sense, however, adaptation appears to imply advance planning, or design, which is misleading.

Abbé refractometer See REFRACTOMETER.

abiogenesis Development of living organisms from non-living matter; as in the supposed origin of life on Earth, or in the concept of spontaneous generation, which was once held to account for the origin of life but which modern understanding of evolutionary processes (see EVOLUTION) has rendered outdated.

abiotic Non-living; devoid of life. Compare BIOTIC.

ablation 1. Removal of snow and ice by melting and by direct alteration from the solid to the gaseous phase (sublimation). The rate of loss is controlled chiefly by air temper-

ature, wind velocity, *humidity, rainfall, and *solar radiation. Ablation on snowfields is also influenced by aspect, depth of snow, and the nature of the underlying surface. Ablation *till is the glacial debris that may be released. The ablation zone of a glacier is that area in which losses, including *calving, exceed additions. 2. Removal of *rock material, especially by wind action.

ablation till See ABLATION 1; and TILL.

ablation zone See ABLATION.

aboral Away from the mouth; on the opposite side of the body from the mouth.

abrasion (corrosion) The erosive (see EROSION) action that occurs when *rock particles of varying size are dragged over or hurled against a surface. Some common agents of abrasion are the *bed load of streams, rock debris embedded in the bases of *glaciers, and *sand and *shingle transported by wind or waves.

abrasion ramp A gentle, seaward slope, with a gradient of approximately 1°, in an intertidal *shore platform that is caused by wave *abrasion. The removal of material by wave action leaves the base of the cliff exposed, leading to further cliff retreat. Abrasion ramps usually terminate on the seaward side where the sea depth reaches about 10 m.

absolute age (true age) The age of a geologic phenomenon measured in present Earth years, rather than its age relative to other geologic phenomena (compare RELATIVE AGE). The term 'absolute age' has been considered rather misleading, as the means for measuring ages (*radiometric dating, *dendrochronology, *varve analysis) are subject to experimental error and the dates obtained are not precise. The alternative term 'apparent age' has been suggested. See also DATING METHODS; and GEOCHRONOLOGY.

absolute humidity See HUMIDITY.

absolute plate motion The motion of a lithospheric *plate (see LITHOSPHERE) with

a

respect to a fixed frame of reference. Various frames of reference have been used, including those defined by *hot spots, no net torque of all the plates, and palaeomagnetic (see PALAEOMAGNETISM) Euler poles (see POLE OF ROTATION).

absolute pollen frequency (APF) *Pollen data from sediments, expressed in terms of the absolute numbers for each *species, *genus, or *family, per unit volume of sediment and, where deposition rates are known, per unit time. In certain circumstances this approach gives clearer information than does the traditional way of expressing pollen data as *relative pollen frequencies (RPFs). APFs are particularly useful in site comparisons in which one or more high pollen producers vary. For example, when trees first appear in the regional pollen rain their prolific pollen may, in an RPF method, give the impression of declining herbaceous species, whereas examination by an APF method will show constant values for herb species.

absolute porosity See POROSITY.

absolute temperature Temperature measured using the *Kelvin scale.

absolute vorticity See VORTICITY.

absolute zero See KELVIN SCALE.

absorptance The ability of a material to absorb *electromagnetic radiation of a specified wavelength. See also ABSORPTANCE BAND.

absorptance band The range of wavelengths of *electromagnetic radiation which are absorbed by a material. See also ABSORPTANCE.

absorption The amount of seismic energy lost during transmission, by conversion to heat. The absorption coefficient is the fractional loss of energy over a distance of one *wavelength; hence higher-*frequency signals are attenuated more readily than those of lower frequencies over the same path. Typical values for *rocks range from 0.25 to 0.75 dB per wavelength.

abstraction (extraction) The artificial removal of water from a well, *reservoir, or river.

Abukama-type metamorphism The *recrystallization of *rocks under a high *geothermal gradient so that at any given temperature the pressure is relatively low. The

term originally referred to a belt of *metamorphic rocks stretching south-westwards from the Abukama Plateau in Japan, and characterized by the development of *andalusite and *sillimanite in rocks that were originally *shales (*pelites). This belt lies parallel to, and on the continental side of, a high-pressure metamorphic belt.

abundance zone See ACME ZONE.

ABW See ARCTIC BOTTOM WATER.

abyssal hills Relatively small topographic features of a dominantly flat, deep-ocean floor, commonly 50–250 m in height and a few kilometres in width. They are most typical of the *Pacific Ocean floor at depths of 3 000–6 000 m.

abyssal plain Smooth, almost level area of the deep-ocean floor in which the gradient is likely to be as low as 1 : 10 000. The covering sediments are usually thin deposits of a *pelagic ooze or *distal *turbidite.

abyssal storm (benthic storm) A large pulse of energy, possibly transferred from the surface, that accelerates *contour currents on the ocean floor to about 40 cm/s, raising large amounts of fine sediment.

abyssal zone Zone of greatest ocean depth, i.e. below a depth of 2 000 m. This zone lies seaward of, and deeper than, the *bathyal zone, and covers approximately 75% of the total ocean floor. It is the most extensive Earth environment, cold, dark, with slow-moving currents (less than a few centimetres per second), supporting *fauna that typically are black or grey, delicately structured, and not streamlined.

Acadian orogeny A phase of mountain building affecting an area from the northern Appalachians in what is now New York State to the Bay of Fundy in maritime Canada (the name refers to the colony of Acadie in that region of French Canada). It occurred in the *Devonian about 390 Ma ago, although the precise date and duration are uncertain, and was most intense east of the Taconic area (see TACONIC OROGENY). It was caused by the westward movement of the Avalon *terrane. See APPALACHIAN OROGENIC BELT.

Acado-Baltic Province See ATLANTIC PROVINCE.

acanthodians See ACANTHODII.

Acanthodii (acanthodians) Class of primitive, fossil fish, characterized by the presence of a true bony skeleton (see BONE), a *heterocercal tail *fin, a persistent *notochord, *ganoid scales, and stout spines in front of the fins. The acanthodians lived from the *Silurian to the *Permian Period and may be related to ancestors of the more modern bony fish.

Acanthograptidae See DENDROIDEA.

Acanthostega See ICHTHYOSTEGA.

acceleration *Evolution that occurs by increasing the rate of ontogenetic (see ONTOGENY) development, so that further stages can be added before growth is completed. This form of *heterochrony was proposed by E. H. Haeckel as one of the principal modes of evolution.

acceleration, gravitational See GRAVITATIONAL ACCELERATION.

accelerograph (earthquake seismometer)

An instrument used to measure *earthquake movements that are too strong for more sensitive *seismometers to register accurately. An accelerograph contains three *accelerometer heads aligned to measure movement in three directions. The accelerograph is often connected directly to the Internet.

accelerometer A device whose output is directly proportional to acceleration. Accelerometers are used in the measurement of the motion of a ship, helicopter, or aircraft during *gravity surveys. A *seismometer or moving-coil *geophone can also function as an accelerometer.

accessory, lithic See LITHIC FRAGMENT.

accessory cloud A small cloud that is seen to be associated with a much larger cloud belonging to one of the ten cloud genera (see CLOUD CLASSIFICATION). *Pileus, *tuba, and *velum are accessory clouds.

accessory mineral A *mineral *phase within a rock whose presence does not affect the root name of the rock. For instance, the root name 'granite' is defined by the presence of *quartz, *alkali feldspar, and *mica. These are the '*essential minerals'. The presence of the mineral *sphene does not affect the root name and hence would be an example of an accessory mineral. *Apatite and *zircon are also common accessory minerals.

accessory plate (sensitive tint) In optical microscopy, a plate used to determine the

optical properties of *minerals. *Quartz, *mica, and *gypsum are the common minerals used to determine the slow and fast *vibration directions that relate to the two *refractive indices of an *anisotropic mineral. The terms 'length-fast' and 'length-slow' may then be assigned to a given mineral for identification purposes. A wedge of quartz (quartz wedge) is used to determine the order of *interference colour exhibited by a mineral.

accidental lithic See LITHIC FRAGMENT.

accommodation space The space in which sediment may accumulate.

accommodation zone A region of intermeshed *normal faults, with very complex geometry, that lies between the boundary faults of a series of interlinked half-grabens (see GRABEN) in an area of crustal extension.

accordion fold See CHEVRON FOLD.

accretion 1. Process by which an inorganic body grows in size by the addition of new particles to its exterior. It is the mechanism by which primitive planetary bodies are believed to form as a result of the accumulation of minute, cold, homogeneous particles (homogeneous accretion). An alternative hypothesis is that iron-rich cores accumulated first and were later surrounded by silicate material (heterogeneous accretion). Homogeneous accretion yields a planet that initially has the same composition from centre to surface; heterogeneous accretion yields a planet that has a layered structure from the start.

2. The accumulation of sediments from any cause, representing an excess of deposition over *erosion.

3. The addition of continental material to a pre-existing continent, usually at its edge. The use of 'accretion' in this sense has evolved from theories of *nucleation to newer theories of the horizontal addition of *allochthonous *terrane of initially coherent bodies of continental *rock, usually more than 100 km² in area, which can collide, rotate, and fragment as they become sutured to a continent.

accretional heating The heating of bodies orbiting a star due to bombardment by smaller objects, the kinetic energy of the impacting body ($\frac{1}{2}mv^2$, where m is mass and v velocity) being released mainly as heat.

accretionary basin A small basin, much smaller than a *fore-arc basin, that develops on the top of an *accretionary wedge and fills

a

with *debris-flow material, *turbidites, and volcanic rocks.

accretionary lapilli Pellets of *ash, ranging in size from 2 mm to 64 mm, which commonly exhibit a concentric ('onion skin') internal structure. The *lapilli are formed by the accretion of very fine ash around condensing water droplets or solid particles, particularly in steam-rich eruptive columns (see ERUPTION). Once formed they can be transported and deposited by *pyroclastic fall, *surge, or flow processes.

accretionary levée See LAVA LEVÉE.

accretionary prism See ACCRETIONARY WEDGE.

accretionary wedge (accretionary prism) A tectonically thickened wedge of *sediment found on the landward side of some *trenches. The accretionary wedge consists of oceanic sediment scraped off the subducting *plate (see SUBDUCTION), plus sediment derived from landward and deposited in the trench. Slices of sediment are added to the wedge by *underthrusting and the trench migrates seaward, the continuation of this process producing an *inversion.

accumulated temperature Surplus or deficit of temperature with respect to a defined mean value and expressed as an accumulation over a given period, e.g. a month, season, or year. For example, a datum value of 6 °C is used as a critical temperature for sustained vegetation growth, against which accumulated surpluses or deficits may be measured.

accumulation zone That part of a *glacier where the mean annual gain of *ice, *firn, and

snow is greater than the mean annual loss. The zone consists of stratified firn and snow together with ice from frozen meltwater. Its lower boundary is the *equilibrium line.


ACD See ARAGONITE COMPENSATION DEPTH.

ACF See ACF DIAGRAM; and AUTOCORRELATION.

ACF diagram A three-component, triangular graph used to show how metamorphic *mineral assemblages vary as a function of *rock composition within one *metamorphic facies. Besides SiO₂, the five most abundant oxides found in *metamorphic rocks are Al₂O₃, CaO, FeO, MgO, and K₂O. The three components plotted on ACF diagrams are A (Al₂O₃), C (CaO), and F (FeO + MgO), making the diagrams particularly useful for showing assemblage variations in metamorphosed, *basic, *igneous rocks, and impure *limestones. However, each of these components has to be modified slightly to account for the presence of other, minor components in the rock. Such modification leads to: A (Al₂O₃ - Na₂O - K₂O); C (CaO - [(10/3)P₂O₅] - CO₂); and F (FeO + MgO - Fe₂O₃ - TiO₂). The minerals *quartz and *albite are assumed to be present in the rocks and are not shown on the diagram. *Tielines connect minerals which coexist in equilibrium and can thus define triangular areas in which three minerals are in equilibrium in the rock, lines on which two minerals are in equilibrium in the rock, and points at which one mineral is in equilibrium in the rock (in addition to the ubiquitous quartz and albite). See AFM DIAGRAM.

achnelith See PELÉ'S HAIR.

achondrite Rare stony *meteorite lacking *chondrules and with low nickel-iron



Accretionary wedge

content. It is more coarsely crystalline than a *chondrite. Basaltic chondrites resemble terrestrial *lavas.

achromatic line In the three-dimensional graph which plots quantities of the three *additive primary colours contributing to *pixels against each other, the line which runs at 45° to the axes. Pixels which plot close to this line will not be strongly coloured and may be subject to *decorrelation stretching.

acicular Pointed or needle-shaped.

acid According to the Brønsted–Lowry theory, a substance that in solution liberates hydrogen *ions or protons. The Lewis theory states that it is a substance that acts as an electron-pair acceptor. An acid reacts with a *base to give a salt and water (neutralization), and has a *pH of less than 7.0. The theory was proposed in 1923 by the Danish physical chemist Johannes Nicolaus Brønsted and the British chemist Thomas Lowry, and independently by the American theoretical chemist Gilbert Newton Lewis.

acidophile An *extremophile (domain *Archaea) that thrives in environments where the *pH is below 5.0.

acid rain Precipitation with a *pH of less than about 5.0, which is the value produced when naturally occurring carbon dioxide, sulphate, and nitrogen oxides dissolve into cloud droplets. The effects of increased acidity on surface waters, soils, and vegetation are complex.

acid rock *Igneous rock containing more than about 60% *silica (SiO₂) by weight, most of the silica being in the form of silicate minerals, but with the excess of about 10% as free *quartz. Typical acid rocks are *granites, *granodiorites, and *rhyolites. *Compare* BASIC ROCK; and INTERMEDIATE ROCK. *See also* ALKALINE ROCK.

acid soil *Soil having a *pH less than 7.0. Degrees of soil acidity are recognized. Soil is regarded as 'very acid' when the reaction is less than pH 5.0. The *USDA lists five standard ranges of soil acidity (less than pH 4.5, extremely acid; 4.5–5.0, very strongly acid; 5.1–5.5, strongly acid; 5.6–6.0, medium acid; and 6.1–6.5, slightly acid). Surface *soil horizons of acid *brown earths have a reaction of pH 5.0 or less.

acme zone (peak zone, flood zone, epibole, abundance zone) An *informal term for a

body of *strata containing the maximum abundance of a particular *taxon occurring within the stratigraphic range of that taxon, and after which the *zone is named.

acoustic impedance (Z) The product of density (ρ) and the acoustic velocity (v) for a given rock mass; $Z = \rho v$. The *reflection coefficient for an interface is governed by the contrast in the acoustic impedances of the two adjacent *rock masses.

acquired characteristics Characteristics that are acquired in the lifetime of an organism, according to early evolutionary theorists such as *Lamarck. Lamarck further suggested that traits acquired in one generation in response to environmental stimuli would be inherited by the next generation. Thus over several generations a particular type of organism would become better adapted (*see* ADAPTATION) to its environment. The kinds of acquisition envisaged by Lamarck and their heritability are now discredited, although there has been a recent revival of some aspects of Lamarckism in modified form.

Acrisols A reference soil group in the soil classification scheme used by the *FAO. Acrisols are *acid soils with an argic B horizon (*see* ARGIC HORIZON) having a *cation-exchange capacity of less than 24 cmol_c/kg.

acritarchs Hollow fossil structures, 5–240 μ m in diameter, inside which dinoflagellates and single-celled algae survived dry periods. They range from *Precambrian to *Recent times. They are found in marine strata, although some non-marine examples are reported from Recent beds. Acritarchs are used in *correlation and to distinguish onshore from offshore *sediments.

Acrothoracica *See* CIRRIPIEDIA.

acrozone *See* RANGE ZONE.

actinium series *See* DECAY SERIES.

actinolite A member of the *amphiboles, Ca₂(Mg,Fe)₃(Si₄O₁₁)₂(OH,F)₂, with the ratio Fe/Fe + Mg = 0.9 to 0.5, belonging to the *tremolite–*ferroactinolite series of Ca-rich amphiboles; sp. gr. 3.0–3.4; *hardness 5–6; *monoclinic; light greenish-grey to dark green; white *streak; *vitreous *lustre; habit *acicular, often fibrous and felted; *cleavage *prismatic, good {110}; occurs widely in low-to medium-grade *schists and some *igneous rocks. The asbestiform variety is called

a

*nephrite and such felted forms were used in the past for insulation and fire-resistant materials, but the development of asbestosis in workers has severely restricted their use.

Actinopterygii (ray-finned fish) A subclass of the *Osteichthyes (bony fish, see BONE), comprising the ray-finned fish, which include the majority of living bony fish of sea and fresh water. The *fins are composed of a membranous web of skin supported by a varying number of spines and soft rays. They appeared first during the *Devonian.

activation analysis See NEUTRON ACTIVATION ANALYSIS.

activation energy (energy of activation) The energy that must be delivered to a system in order to increase the incidence within it of reactive molecules, thus initiating a reaction.

active geophysical methods Geophysical exploration methods which require an artificial signal to be generated. For example, exploration seismology, some *electromagnetic techniques, *electrical resistivity, *remote sensing, and *induced polarization are said to be active geophysical methods. The term is contrasted with *passive geophysical methods.

active layer Seasonally thawed surface layer between a few centimetres and about 3m thick, lying above the permanently frozen ground in a periglacial environment. It may be subject to considerable expansion on freezing, especially if silt-sized particles dominate, with important engineering implications. See also MOLLISOLS; and PERMAFROST.

active margin (seismic margin) The margin of a continent that is also a *plate margin. The alternative term, 'Pacific-type margin', indicates the range of features (e.g. *earthquakes, andesitic (see ANDESITE) volcanic chains, offshore oceanic *trenches, and young fold mountains) which may be associated with active margins. Some authors distinguish an 'Andino-type margin', involving an oceanic and a continental plate, from a 'Japan-type margin', involving an oceanic plate and an *island arc. The term 'Mediterranean-type margin' is also in use, although to a lesser extent, to signify the coincidence of continental edges and plate margins in a *collision zone.

active methods See ACTIVE GEOPHYSICAL METHODS.

active pool The part of a *biogeochemical cycle in which the nutrient element under consideration exchanges rapidly between the biotic and abiotic components. Usually the active pool is smaller than the *reservoir pool, and it is sometimes referred to as the 'exchange' or 'cycling' pool.

active remote sensing *Remote sensing which is based on the illumination of a scene by use of artificial radiation. An example is *radar. Compare PASSIVE REMOTE SENSING.

activity A broadly used term which refers to the rate or extent of a change associated with some substance or system. For example, it may be the tendency of a metal high in the electromotive series to replace another metal lower in the series, e.g. magnesium displacing copper from most of its compounds. It may also be used to describe the rate of decay of atoms by radioactivity.

activity coefficient (γ) The ratio of chemical activity (i.e. the effective concentration, a) of a component in a solution, to the actual mole fraction (X) present in solution: ($\gamma = a/X$). Values for activities are determined experimentally in a number of ways, including measuring the ratio of the *vapour pressure (p) of a known concentration of the substance in solution to the vapour pressure (p^*) of the pure substance: $a = p/p^*$. In an ideal solution the activity coefficient = 1, and the activity of the component is equal to its mole fraction. In general, the greater the amount of dissolved material, the lower the activity coefficients of each of the species present.

Actonian A *stage of the *Ordovician (453–454 Ma ago) in the Upper *Caradoc, underlain by the *Marshbrookian and overlain by the *Onnian.

actual evapotranspiration (AE) The amount of water that evaporates from the surface and is transpired by plants if the total amount of water is limited. Compare POTENTIAL EVAPOTRANSPIRATION.

actualism The theory that present-day processes provide a sufficient explanation for past geomorphological phenomena, although the rate of activity of these processes may have varied. The theory was first clearly expressed in 1749 by G. L. L. *Buffon (1707–88), and was the essential principle of *uniformitarianism as presented in 1830 by C. *Lyell (1797–1875).

acuity The ability of a human to discern spatial variation in a scene.

ACV See AGGREGATE TESTS.

Adam The postulated male ancestor for all modern humans, who lived in Africa between about 100 000 and 200 000 years ago. 'Adam' is based on a change in the human Y chromosome that occurred at that time in one descendant of Adam and is now present in all human males, except for some Africans. See also MITOCHONDRIAL EVE.

adamantine Of mineral *lustre, brilliant, like a polished diamond.

adamellite A rock of granitic composition (see GRANITE) characterized by the presence of *quartz, *plagioclase feldspar, and potassic feldspar (see ALKALI FELDSPAR) accompanied by *biotite and/or *hornblende. The two feldspar types occur in approximately equal proportions, the plagioclase composition lying within the oligoclase range. The name is derived from the type locality of Adamello in the Tyrol where granites of this type were originally defined. In Britain the best-known example occurs at Shap Fell in Cumbria.

Adams–Williamson equation Equation describing a fundamental relationship between seismic velocities (v_p and v_w), the *gravitational acceleration (g), and the adiabatic change in density (dp) within the *Earth (assuming only hydrostatic pressure) as a function of radius (dr):

$$dp = \frac{gp}{drv_p^2 - (4/3)v_w^2}$$

This equation is directly applicable to the lower *mantle and outer *core, but is invalid where the composition is variable, the pressure is not hydrostatic, or the increase in pressure is not adiabatic.

adapical A directional term meaning towards the shell *apex.

adaptation 1. Generally, the adjustments that occur in animals in respect of their environments. The adjustments may occur by *natural selection, as individuals with favourable genetic traits breed more prolifically than those lacking these traits (genotypic adaptation), or they may involve non-genetic changes in individuals, such as physiological modification (e.g. acclimatization) or behavioural changes (phenotypic adaptation). Compare ABAPTATION. **2.** In an evolutionary sense, that which fits an organ-

ism both generally and specifically to exploit a given environmental zone.

adaptive radiation 1. A burst of evolution, with rapid divergence from a single ancestral form, resulting in the exploitation of an array of habitats. The term is applied at many *taxonomic levels, e.g. the radiation of the mammals at the base of the *Cenozoic refers to *orders, whereas the radiation of 'Darwin's finches' in the Galápagos Islands resulted in a proliferation of *species. **2.** Term used synonymously with *cladogenesis by some authors.

adaptive zone The adaptive specialization(s) that fit the *taxon to its environment, e.g. feeding habits.

addition rule (Weiss zone law) With reference to crystallographic notation, the rule stating that the indices (see MILLER INDEXES) of two *crystal faces in the same *zone always add up to the indices of a face bevelling the edge lying between them. The rule may be used to index faces on a *stereogram, or faces at the intersection of two zones.

additive primary colours The spectral colours red, green, and blue, which, when mixed together by projection through filters, can be used to produce all other colours. None of the primary colours can be produced by combinations of the other two. See also SUBTRACTIVE PRIMARY COLOURS.

adductor muscles See MUSCLE SCAR.

Adelaidean A *stage (542–1 300 Ma ago) of the Upper *Proterozoic of south-eastern Australia, underlain by the *Carpenterian and overlain by the Hawker (*Cambrian).

Adelaidean orogeny A late *Proterozoic and *Ordovician phase of mountain building, affecting what is now southern Australia, in which *sedimentary rocks of the Adelaide System were raised by severe thrusting and overfolding, first in the south and later along the northern margin of the system.

adhesion ripples See ADHESION WARTS.

adhesion warts (adhesion ripples) A *sedimentary structure consisting of an irregular, wart-like or blistered, *sand surface, formed by the wind blowing dry sand over a moist surface. The warts tend to be slightly asymmetrical, with steeper sides in the upwind direction.

a

adiabat The rate at which a *parcel of air cools as it rises and warms as it descends, as indicated by two lines (dry adiabat and wet adiabat) on a *tephigram.

adiabatic Applied to the changes in temperature, pressure, and volume in a *parcel of air or liquid that occur as a consequence of the vertical movement of the fluid, and without any exchange of energy with the surrounding fluid. *See also* DRY ADIABATIC LAPSE RATE; and SATURATED ADIABATIC LAPSE RATE.

adit Horizontal or nearly horizontal tunnel from the surface into a mine, for entry, drainage, or exploration.

admission The substitution of a *trace element for a major element with a similar *ionic radius but a higher *valency during the crystallization of a *magma, e.g. the substitution of Li^+ for Mg^{2+} in the *pyroxenes, *amphiboles, and *micas.

adobe A silty *clay, often calcareous, found in dry, desert-lake basins. This fine-grained *sediment is usually deposited by desert floods which have eroded wind-blown *loess deposits. The term is of Spanish origin.

adoral On the same side of the body as the mouth.

Adrastea (Jupiter XV) A jovian satellite (a *moon) that orbits within the main ring of Jupiter; it and *Metis may be the source of the material comprising the ring. Both are considered too small to suffer tidal disruption, but eventually their orbits will decay. Adrastea is one of the smallest satellites in the solar system. It was discovered in 1979 by David Jewitt. Its diameter is 20 km (± 20) ($23 \times 20 \times 15$ km); mass 1.91×10^6 kg; mean distance from Jupiter 129 000 km.

adsorption The attachment of an ion, molecule, or compound to the charged surface of a particle, usually of *clay or *humus, from where it may be subsequently replaced or exchanged. Ions carrying positive charges (e.g. those of calcium, magnesium, sodium, and potassium) become attached to, or adsorbed by, negatively charged surfaces (e.g. those of clay or humus).

adsorption complex Various materials of the soil, mainly *clay and *humus and to a lesser degree other particles, capable of adsorbing ions and molecules.

adularia *See* ALKALI FELDSPAR.

advection The horizontal transfer of heat by means of a moving gas (usually air).

adventive cone *See* PARASITIC CONE.

AE *See* ACTUAL EVAPOTRANSPIRATION.

aedifichnia A category of *trace fossils that comprises structures in full relief that were constructed by organisms from raw materials, e.g. mud nests of wasps, caddis fly cases, spiders' larders consisting of concentrations of insects, insect remains, and spiders.

aegirine *Pyroxene mineral, $\text{NaFe}^3\text{Si}_2\text{O}_6$; sp. gr. 3.5; *hardness 6; *monoclinic; greenish-black or brown; occurs as fairly short, *prismatic crystals in *igneous and *metamorphic rocks. A variety intermediate in composition between aegirine and augite is called 'aegirine-augite'. *See also* AUGITE; and CLINOPYROXENE.

Aegyptopithecus zeuxis A genus and species of early *catarrhine primates, known from abundant remains, including several nearly complete skulls, from the early *Oligocene of the Jebel al-Qatrani Formation, Fayum, Egypt. The size of a small, living monkey, it had a long tail and could jump from branch to branch. It possessed the dental and some of the cranial characteristics of living catarrhines, but lacked many of the other cranial and most of the postcranial diagnostic features, and so represents a time when catarrhines had separated from other primates, but remained more primitive than living hominoids (*Hominoidea) or Old World monkeys and it could have been ancestral to living catarrhines.

aeolian abrasion The erosion of a surface that is caused by bombardment with loose particles carried by the wind.

aeolianite General term for the sedimentary products of wind (aeolian) deposition.

aeolian processes (eolian processes) The erosion, transport, and deposition of material due to the action of the wind at or near the Earth's surface. Aeolian processes are at their most effective when the vegetation cover is discontinuous or absent.

aeolian ripple (eolian ripple) A ripple on the surface of a sedimentary rock that is caused by saltating grains. Aeolian ripples have a wavelength approximately equal to the *saltation path of the grains, and they usually have no internal *cross-lamination.

Aeolis Quadrangle A region of Mars formed in the Late *Noachian or Early *Hesperian Epoch, containing both extensional and compressional land-forms and *valles, some of which may be outflow channels, but some of which may be tectonic rift features.

aerial photograph A photograph taken from an aircraft. In hydrology, false-colour infrared photographs are used to determine the wetness and temperature of soils and to detect *springs.

aerial photography The taking of aerial photographs of rock exposures and of the ground surface for purposes of geologic interpretation. The photographs may be taken vertically, or at a high-oblique or low-oblique angle, and may be assembled like a mosaic to provide a picture of a large area. Stereoscopic cameras (two cameras within a single body) may be used to produce pairs of pictures that provide three-dimensional pictures when observed through a stereoscopic viewer. See PHOTO GEOLOGY.

aerobic 1. Of an environment: one in which air (oxygen) is present. In the case of a depositional environment, one with more than 1 ml of dissolved oxygen per litre of water. Compare ANAEROBIC; and DYSAEROBIC. **2.** Of an organism: one requiring the presence of oxygen for growth, i.e. an aerobe. **3.** Of a process: one that occurs only in the presence of oxygen.

aerodynamic roughness Uneven flow of air caused by irregularities in the surface (which may be of a solid, or of air of different density) over which the flow takes place.

aerological diagram Diagram to demonstrate variations with height of the physical characteristics of the atmosphere, particularly its temperature, pressure, and *humidity.

aeromagnetic survey Survey of the Earth's magnetic field, based on data from *magnetometers towed behind aircraft or suspended below helicopters. These instruments measure the total intensity of the *geomagnetic field or, occasionally, components of this field. The resulting measurements can then be compared with theoretical models for the value of the field and the differences (*magnetic anomalies) can be interpreted in terms of changes in the magnetic properties of the rocks below the survey line or grid. The magnetometers are usually flown with other instrumentation, e.g. *radiometric and electromagnetic, at the lowest practicable con-

stant height above the ground. Usually the magnetometer is housed in a 'bird' towed behind the aircraft, or in a wing-tip pod, or in a 'stinger' in the tail. In cases where the magnetometer is on board, in-board coil systems compensate for the aircraft's own magnetic field.

Aeronian A *stage (436–439 Ma ago) of the Lower *Silurian (*Llandovery Period) underlain by the *Rhuddanian and overlain by the *Telychian.

aerosol Colloidal substance, either natural or man-made, that is suspended in the air because the small size (0.01–10 μm) of its particles makes them fall slowly. Aerosols in the *troposphere are usually removed by *precipitation and their *residence time is measured in days or weeks. Aerosols that are carried into the *stratosphere usually remain there much longer. Tropospheric aerosols may act as *Aitken nuclei but the general effect of aerosols is to absorb, reflect, or scatter radiation. Stratospheric aerosols, mainly sulphate particles resulting from volcanic *eruptions, may reduce *insolation significantly. About 30% of tropospheric dust particles are the result of human activities. See ATMOSPHERIC STRUCTURE; MIE SCATTERING; RAYLEIGH SCATTERING; and VOLCANIC DUST.

Aëtosauria Mainly *Triassic group of primitive thecodontian ('tooth-in-socket') reptiles (see THECODONTIA). They resembled heavily armoured crocodiles, and appear to have been specialized herbivores or possibly omnivores. They grew up to 3 m long, and their armour plating comprised rows of bony *plates.

AFc See ASSIMILATION-FRACTIONAL CRYSTALLIZATION.

AF demagnetization See ALTERNATING MAGNETIC FIELD DEMAGNETIZATION.

AFM diagram A three-component, triangular graph used to show how metamorphic *mineral assemblages vary as a function of *rock composition within one *metamorphic facies. Besides SiO₂, the five most abundant oxides found in *metamorphic rocks are Al₂O₃, CaO, FeO, MgO, and K₂O. The three components plotted on AFM diagrams are derived from a tetragonal diagram, with species Al₂O₃, K₂O, FeO, and MgO, and are ideal for showing mineral assemblage variations as a function of the composition of *pelites. Mineral and rock compositions plotting within this diagram are projected on to the Al₂O₃-FeO-MgO face from either the *muscovite or K-feldspar point on

a

the Al_2O_3 -FeO edge. The components of the diagram are thus A (Al_2O_3), F (FeO), and M (MgO), with the projection geometry being accommodated on specially scaled axes. Each of these components has to be modified slightly to account for the presence of other, minor components in the rock, leading to: A ($\text{Al}_2\text{O}_3 - 3\text{K}_2\text{O}$); F ($\text{FeO} - \text{TiO}_2 - \text{Fe}_2\text{O}_3$); and M (MgO). The minerals *quartz and *albite are assumed to be present in the rocks and are not shown on the diagram. As in *ACF diagrams, *tielines connect minerals which coexist in equilibrium.

AFMAG EM system Audio-Frequency Magnetic ElectroMagnetic method, which uses natural electromagnetic (EM) fields (*sferics) in the audio-frequency range (1–1000 Hz) generated by thunderstorms to investigate lateral changes in the *resistivity of the Earth's surface.

African Plate One of the present-day major lithospheric *plates, consisting of the continental mass of Africa surrounded, except to the north, by *oceanic crust and oceanic *ridges. To the north, a complex picture of collision and *subduction zones and *transform faults has been postulated for the boundary with the *Eurasian Plate and various minor plates, e.g. the *Aegean Plate. The northern part of the African Plate also contains remnants of the oceanic crust of *Tethys. To the north-east the Red Sea is interpreted as an actively forming ocean, at the young stage of the *Wilson cycle, while the E. African *rifts, partially defining what is called by some the 'Somali Plate' to the east, may be at the embryonic stage of ocean development, or possibly a stillborn ocean.

aftershock A seismic event that occurs after an *earthquake, usually within days or weeks. Although often of small *magnitude, aftershocks can be more destructive as buildings and structures have already been weakened.

Aftonian The earliest (1.3–0.9 Ma) of four *interglacial *stages in N. America, following the *Nebraskan glacial episode, and approximately equivalent to the *Donau/Günz interglacial of Alpine terminology. Climatically it was marked by mild summers and winters warmer than those in present-day N. America.

Agassiz, Jean Louis Rodolphe (1807–73) A Swiss geologist who worked initially on fossil fish, Agassiz is better known for his *glacial theory (1837). He met *Buckland in 1840, and persuaded him that *drift deposits

in Britain were evidence of a glacial epoch. In 1846 he moved to the USA to become professor of zoology and geology at Harvard, where he founded the Museum of Comparative Zoology (1859).

agate (mocha stone) Variety of chalcidonic silica (SiO_2) that is *cryptocrystalline. It is similar to *chalcedony except that impurities of iron and manganese may give it a distinct colour banding which is frequently precipitated in concentric zones. Moss agate contains delicate, fern-like, dendritic patterns. Agates may be cut and polished as decorative stones.

age 1. The interval of geologic time equivalent to the *chronostratigraphic unit '*stage'. Ages are subdivisions of *epochs and may themselves be subdivided into *chrons. An age takes its name from the corresponding stage, so like the stage name it carries the suffix '-ian' (or sometimes '-an'); the term 'age' is capitalized when used in this formal sense, e.g. '*Oxfordian Age'. **2.** An *informal term to denote a time span marked by some specific feature, e.g. '*Villefranchian mammalian age'.

ageostrophic wind The vector difference between the *geostrophic and the actual winds.

agglomerate Coarse-grained volcanic rock with rounded to subangular fragments. These fragments are mainly larger than 2 cm in size, but the mixture of fragments is typically ill sorted and the *matrix may be fine grained. An agglomerate may be the product of a volcanic explosion and therefore a *pyroclastic rock, but often the term 'agglomerate' is applied to brecciated volcanic rocks of uncertain origin. Those deposits may range from vent *breccias to debris from mudflow or lahar deposits.

agglutinate A constituent of lunar soils comprising glass-bonded *aggregates, which consist of *glasses and rock and mineral fragments welded together by glass. These aggregates form during the impact of micrometeorites into lunar soils. Their abundance in a lunar soil is an index of exposure to micrometeorite bombardment, and hence to soil maturity. The average size of agglutinates in mature soils varies, but tends toward a mean of 60 μm .

aggradation The general accumulation of unconsolidated sediments on a surface, which thereby raise its level. A large range of

mechanisms may be involved, including *fluvial, *aeolian, marine, and *slope processes.

aggregate 1. In the building and construction industry, a mixture of mineral substances (bulk *minerals), e.g. sand, gravel, crushed *rock, stone, slag, and other materials (e.g. colliery spoil, pulverized fuel ash) which, when cemented, forms *concrete, mastic, mortar, plaster, etc. Uncemented, it can be used as road-making material, railway ballasts, filter beds, and in some manufacturing processes as flux. In road-making, aggregate mixed with *bitumen is called 'coated stone', and different physical characteristics are required for the different layers comprising the road *pavement. Fine aggregate is less than 6.35 mm in diameter, coarse aggregate greater than 6.35 mm. See AGGREGATE TESTS; and PAVEMENT.

2. Group of soil particles adhering together in a cluster; the smallest structural unit, or ped, of soil. Aggregates join together to make up the major structural soil units.

aggregate abrasion value See AGGREGATE TESTS.

aggregate crushing value See AGGREGATE TESTS.

aggregate impact value See AGGREGATE TESTS.

aggregate tests Specific tests used to determine the suitability of *aggregates for special purposes. There are tests for: (a) shape and texture (the angularity number), to determine whether particles have a large angle of friction with good bonding properties; (b) size and grading, to determine whether particles will pack well; (c) moisture content, to discover whether materials absorb so much water that freeze-thaw action might cause the break-up of structures; (d) rock density, which may affect the economics of an operation; (e) strength, determined by subjecting the rock to hammering in a standard test and measuring the percentage of fine material produced (the aggregate impact value, or AIV); (f) resistance to crushing (the aggregate crushing value, or ACV), measured in a similar manner; (g) resistance to abrasion, measured by standard equipment to give the aggregate abrasion value (AAV)—the lower the AAV, the more resistant the rock; and (h) resistance to polishing, measured in the laboratory to give the polished stone value (PSV)—the higher the PSV, the greater the resistance to polishing

and therefore skidding, and the more valuable the material.

aggregation 1. Process in which soil particles coalesce and adhere to form soil aggregates. The process is encouraged by the presence of bonding agents such as organic substances, *clay, iron oxides, and ions (e.g. calcium and magnesium). **2.** Progressive attachment of particles (e.g. ice or snow) or droplets around a nucleus, thereby causing its growth.

Aglaophyton major See RHYNIA.

Agnatha (phylum *Chordata, subphylum *Vertebrata) Superclass of jawless, fish-like vertebrates, with sucker-like mouths, including the extant lampreys, slime-eels and hagfish, and some of the earliest primitive vertebrates, with heavily armoured forms, e.g. **Cephalaspis* (see also OSTEOSTRACI), *Pteraspis* (see HETEROSTRACI), and *Jamoytius* (see ANASPIDA). They appeared first during the *Ordovician.

Agnostida An order of *Trilobita that lived from the Lower *Cambrian to Upper *Ordovician. Most were blind, lacking sutures, and typically are found rolled up. They had a subequal *cephalon and *pygidium. There were two suborders. They are important stratigraphic markers.

agric horizon Mineral-soil diagnostic horizon formed from an accumulation of *clay, *silt, and *humus, which has moved down from an overlying, cultivated soil layer. It is a *soil horizon created by agricultural management, and is identified by its near-surface position, and by *colloids accumulated in the pores of the soil.

agrichnion (pl. agrichnia) A *trace fossil comprising a burrow that formed the permanent dwelling of an organism and was used to trap or culture smaller organisms for food or use them in chemosymbiosis.

Agricola, Georgius (Georg Bauer) (1494–1555) The author of works on 'geology' and mineral classification, and of the first comprehensive record of mining, *De Re Metallica* (1556). Using Roman sources and contemporary German knowledge, his books became basic reference material for two centuries.

agrometeorology The study of the relationship between conditions in the surface layers of the atmosphere and those in the surface of the Earth, as this affects agriculture.

AGU See AMERICAN GEOPHYSICAL UNION.